

---

# ***Investigating the Influence of Climate Variability and Topographic Factors on Vegetation Dynamics and Habitat Diversity within the Schorfheide-Chorin Biosphere Reserve.***

---

Yourfee Kennedy, Biosphere Reserves Management, MSc. Eberswalde University for Sustainable Development, Germany.

## **Abstract**

This investigation delves into the effects of climate variability and topographic elements on vegetation dynamics and habitat diversity within the Schorfheide-Chorin Biosphere Reserve located in northeastern Germany. By utilizing multi-year Normalized Difference Vegetation Index (NDVI) data obtained from Landsat 8, climate data sourced from the WorldClim database, and topographic details from the USGS Shuttle Radar Topography Mission (SRTM), an analysis of spatial and temporal patterns in vegetation productivity was conducted. The results indicate notable fluctuations in vegetation distributions influenced by changes in climate and variations in terrain, thus emphasizing the intricate relationships among these components. This study highlights the significance of incorporating climate and topographic factors into ecosystem management and conservation approaches to promote biodiversity and enhance resilience within biosphere reserves.

**Keywords:** climate variability, topographic factors, vegetation dynamics, habitat diversity, Schorfheide-Chorin, biosphere reserve, NDVI, remote sensing, Landsat 8, and ecosystem management

# 1 Introduction

Biosphere reserves are designated areas that function as exemplars for the promotion of sustainable development and the conservation of biodiversity. The Schorfheide-Chorin Biosphere Reserve, situated in the northeastern region of Germany, represents a distinctive ecosystem characterized by a variety of landscapes, encompassing vast pine forests, deciduous woodlands, and grasslands. It is imperative to comprehend the dynamics of vegetation and habitat diversity within this reserve to facilitate efficient management and conservation strategies (Hartoyo et al., 2022; Nasiri et al., 2022). The impact of climate variability and topographic elements on the configuration of vegetation patterns and the distribution of habitats is well-documented. Alterations in precipitation and temperature patterns can impact plant development, productivity, and phenological cycles, potentially leading to changes in vegetation communities. Furthermore, topographic attributes like elevation, slope, and aspect can generate microclimatic conditions that influence soil moisture, nutrient availability, and exposure to sunlight, consequently affecting vegetation dynamics (Almalki et al., 2022; Ding et al., 2020; Han et al., 2020; Y. Liu et al., 2021; Pei et al., 2021; Tao et al., 2018; Vicente-Serrano et al., 2010). The primary objective of this research is to explore the impact of climate variability and topographic factors on vegetation dynamics and habitat diversity in the Schorfheide-Chorin Biosphere Reserve. By incorporating multi-year Normalized Difference Vegetation Index (NDVI) data derived from Landsat 8 satellite imagery, climate data sourced from the WorldClim database, and topographic details from the USGS Shuttle Radar Topography Mission (SRTM) dataset, this study aims to offer valuable insights into the spatial and temporal trends of vegetation alterations and their potential driving forces (Dubovyk et al., 2016; He et al., 2023; Hossain & Li, 2021; Johansson et al., 2013; Knapp & Smith, 2001; Li et al., 2023; Lian et al., 2021; Liang et al., 2013; Lin et al., 2017; Q. Liu et al., 2018; Seo & Kim, 2021; Zheng et al., 2021; Zhou et al., 2011).

## 2 Literature Review

Biosphere reserves have been designated by UNESCO as exemplary regions for sustainable development, biodiversity conservation, and ecosystem monitoring amidst conditions of global change. These specific areas function as operational environments for analyzing the interplay between human activities and the environment, advocating for sustainable land utilization practices, and bolstering research and educational endeavors. Remote sensing methodologies, notably the utilization of vegetation indices such as NDVI, have demonstrated their effectiveness as valuable instruments for monitoring vegetation dynamics and evaluating habitat diversity. NDVI serves as a widely utilized metric for gauging vegetation verdancy, productivity, and phenological variations, and has been utilized in diverse research undertakings to scrutinize the repercussions of climate variability and topographic elements on vegetation patterns(Chen et al., 2015; Cong et al., 2013; B. Fu et al., 2014; Y. H. Fu et al., 2021; Mao et al., 2012; Menzel et al., 2006; Rechid et al., 2009; Suepa et al., 2016; Tang et al., 2015; Tu et al., 2018; Yu et al., 2021; P. Zhang et al., 2020; X. Zhang et al., 2004; Y. Zhang et al., 2014).Climate parameters, like precipitation and temperature, wield a pivotal influence in configuring vegetation communities and ecosystem functions(Gado et al., 2019; Jiao et al., 2021). Alterations in precipitation trends can impact soil moisture availability, consequently influencing plant growth and productivity, while fluctuations in temperature can affect phenological rhythms and species distributions. Various studies have showcased the impact of climate variability on vegetation dynamics through the utilization of remote sensing information and climate models(Aslami & Ghorbani, 2018; Islam et al., 2021; Novillo et al., 2019).Topographic elements, encompassing elevation, slope, and aspect, possess the capacity to engender microclimatic circumstances that shape vegetation distributions and habitat diversification.(Ettehadi Osgouei & Kaya, 2017; Nemani et al., 1996).Disparities in

topography can influence soil moisture levels, nutrient accessibility, and exposure to sunlight, which can culminate in unique vegetation assemblages and species compositions. Remote sensing methodologies, when merged with topographic data, have been harnessed to explore the correlations between topographic factors and vegetation dynamics across diverse ecosystems(Aini Rahmi et al., 2024; Camps-Valls et al., 2021; Papagiannopoulou et al., 2017)

### 3 Methodology

The methodology proposed exploits the strengths of remote sensing data, climate information, and topographic data to offer a comprehensive insight into the factors influencing vegetation dynamics and habitat diversity in the Schorfheide-Chorin Biosphere Reserve. The utilization of QGIS as the principal software platform guarantees the accessibility and reproducibility of the analytical workflow(Huang et al., 2021; Lambin et al., 2001; Lu et al., 2004).Data Sources:


1. Normalized Difference Vegetation Index (NDVI) Data: Multi-year NDVI data will be sourced from Landsat 8 satellite imagery via the United States Geological Survey (USGS) website. This data will enable the examination of vegetation dynamics and phenological variations in the study area over the specified timeframe.
2. Climatic Data: Precipitation and temperature data for 2020 and 2021 (Al Shawabkeh et al., 2023)will be obtained from the WorldClim database. These climatic parameters will be utilized to evaluate the impact of climate fluctuations on vegetation patterns within the biosphere reserve.
3. Topographical Data: The Digital Elevation Model (DEM) for the study region will be acquired from the USGS Shuttle Radar Topography Mission (SRTM) dataset. This DEM will be instrumental in deriving topographic features like elevation, slope, and aspect, which will be integrated into the analysis to explore the influence of topographic elements on vegetation dynamics and habitat diversity(Khalile et al., 2018; Poyatos et al., n.d.).

Data Processing and Analysis:


1. Processing of NDVI Data: The Landsat 8 NDVI data will undergo preprocessing in QGIS, encompassing tasks such as atmospheric correction, cloud masking, and mosaicking of scenes to generate a seamless NDVI dataset for the study area.
2. Processing of Climatic Data: The precipitation and temperature data extracted from the WorldClim database will be processed for the study area using QGIS (Matsushita et al., 2007; Rutkoski et al., 2016). Spatial interpolation methods may be utilized to produce continuous climate variable surfaces if needed.
3. Processing of Topographical Data: The SRTM DEM will be processed in QGIS to derive elevation, slope,

and aspect for the study area. 4. Integration and Analysis of Data: The processed NDVI, climate, and topographical data will be integrated within QGIS for analysis(Badamasi et al., n.d.; P. Li et al., 2020; Lioubimtseva & Henebry, 2009; Propastin et al., 2008)

#### 4 Results




**Figure 1. Spatial distribution of vegetation productivity in Schorfheide-Chorin Biosphere Reserve 2020**




**Figure 2. Spatial distribution of vegetation productivity in Schorfheide-Chorin Biosphere Reserve 2021**





**Figure 3. Spatial distribution of vegetation productivity in Schorfheide-Chorin Biosphere Reserve 2022**



**Figure 4. Spatial distribution of vegetation productivity in Schorfheide-Chorin Biosphere Reserve 2023**



0°00' 90°00'E

0°00' 90°00'E

0°00' 90°00'E

0°00' 90°00'E

Schorfheide-Chorin

NDVI Values for Vegetation Cover

2024 NDVI Values

- Moderate Vegetation  $\leq 0.64$
- Dense vegetation  $= 0.64 - 0.68$
- Lush Vegetation cover  $= 0.68 - 0.69$
- Very Dense  $> 0.69$

1:247,406

8

Topographic map of the Schorfheide-Chorin region in Germany. The map displays elevation ranges categorized into four levels: Lowland (cyan, ≤ 55m), Undulating (yellow, 55-67m), Elevated (orange, 67-78m), and Upland (dark grey, > 78m). The map is bounded by 0°00' to 9°00' East and 52°00' to 53°00' North. A black cross symbol is present in the top right corner. The scale is 1:61,726.

9

## Temperature Distribution in the Schorfheide-Chorin Biosphere Reserve in 2020




Figure 7. Temperature distribution of the Schorfheide-Chorin Biosphere Reserve in 2020

## Temperature Distribution in the Schorfheide-Chorin Biosphere Reserve in 2021




Figure 8. Temperature distribution of the Schorfheide-Chorin Biosphere Reserve in 2021

# Spatial Distribution of Annual Precipitation in the Schorfheide-Chorin Biosphere Reserve (2020)




Figure 9. Spatial Distribution of Annual Precipitation 2020

# Spatial Distribution of Annual Precipitation in the Schorfheide-Chorin Biosphere Reserve (2021)




Figure 10. Spatial Distribution of Annual Precipitation 2021

## 5 Discussion

The outcomes of this investigation offer valuable insights into the impact of climate variability and topographic elements on vegetation dynamics and habitat diversity within the Schorfheide-Chorin Biosphere Reserve. The multi-year NDVI data (Figures 1-5) depict clear spatial patterns and temporal fluctuations in vegetation productivity throughout the reserve (de la Fuente & Williams, 2022). Within the year 2020 (Figure 1), the NDVI values varied from moderate ( $\leq 0.61$ ) to luxuriant vegetation ( $> 0.72$ ), indicating a diverse dispersion of vegetation cover and productivity levels. The subsequent year, 2021 (Figure 2), displayed a similar array of NDVI categories, with certain regions showing slightly elevated or diminished productivity compared to the previous year. This inter-annual variability in vegetation productivity may be ascribed to fluctuations in climatic conditions, specifically precipitation, and temperature, as demonstrated by the climate data for 2020 and 2021 (Figures 7-10) (Filippi et al., 2021). The NDVI data for 2022 (Figure 3) unveiled a significant change in vegetation productivity, with the appearance of a sparse vegetation category ( $\leq 0.47$ ) and a greater proportion of areas classified as dense vegetation (0.48-0.5) or lush vegetation ( $> 0.5$ ). This transformation could be associated with particular climatic occurrences or patterns during that year, possibly impacting soil moisture availability, temperature patterns, or other environmental aspects influencing plant growth and productivity (Filippi et al., 2021). In the year 2023 (Figure 4), the NDVI categories were delineated differently, extending from thriving vegetation ( $\leq 0.71$ ) to exceedingly dense vegetation ( $> 0.78$ ). This variability in category definitions underscores the dynamic nature of vegetation productivity and the necessity for adaptive classification systems to encompass the complete spectrum of conditions within the biosphere reserve (Piao et al., 2006). The 2024 NDVI data (Figure 5) exhibited a return to more moderate vegetation productivity levels, with categories stretching from moderate ( $\leq 0.64$ ) to highly dense

vegetation ( $>0.69$ ). This fluctuation in vegetation patterns over the five years emphasizes the impact of climate variability and the potential repercussions of extreme events or irregularities on the ecosystems of the biosphere reserve (Hussain et al., 2023). The topographic depiction (Figure 6) disclosed the varied terrain characteristics within the Schorfheide-Chorin Biosphere Reserve, spanning from lowlands ( $\leq 55$  m) to elevated areas ( $>78$  m). These topographic variances likely play a role in shaping microclimatic conditions and influencing soil moisture availability, nutrient dispersion, and exposure to sunlight, ultimately molding the distribution and composition of vegetation communities within the reserve. (C. Liu et al., 2021) The temperature and precipitation data for 2020 and 2021 (Figures 7-10) provided valuable insights into the climatic conditions that might have impacted vegetation dynamics during those years. The temperature categories fluctuated from sub-zero ( $\leq 2^{\circ}\text{C}$ ) to above-freezing ( $>2.4^{\circ}\text{C}$ ) in 2021, while 2020 showcased a different spectrum, from cold ( $\leq 6^{\circ}\text{C}$ ) to warm ( $>6.1^{\circ}\text{C}$ ) conditions. Similarly, the precipitation data displayed variances in the annual rainfall quantities, with categories spanning from arid ( $\leq 34.9$  mm) to copious ( $>39.9$  mm) in 2020 and from dry ( $\leq 51.2$  mm) to plentiful ( $>58.8$  mm) in 2021. The climatic fluctuations, in conjunction with the diverse topographical landscape, probably played a role in shaping the spatial distributions and temporal fluctuations in vegetation productivity across the biosphere reserve. Regions characterized by greater altitudes or arid climates may have exhibited reduced vegetation productivity, whereas low-lying areas or regions with more optimal levels of moisture and temperature might have sustained denser vegetation (Zheng et al., 2019).

## 6 Conclusion

This research has illustrated the impact of climate variability and topographic elements on vegetation dynamics and habitat diversity within the Schorfheide-Chorin Biosphere Reserve. The multi-year NDVI data has unveiled distinct spatial patterns and temporal variations in vegetation productivity, potentially influenced by fluctuations in precipitation, temperature, and other climatic factors(S. Li et al., 2023)The diverse topography of the reserve, spanning from low-lying areas to highland regions, likely played a role in creating microclimatic conditions and affecting soil moisture levels, nutrient distribution, and exposure to sunlight, thus shaping the distribution and makeup of vegetation communities The amalgamation of remote sensing data (NDVI), climate data (temperature and precipitation), and topographic information (DEM) has offered a comprehensive comprehension of the intricate interactions among these elements and their consequences on vegetation dynamics and habitat diversity within the biosphere reserve(Al-Doski et al., n.d.)The results of this investigation underscore the significance of incorporating both climate variability and topographic aspects in the monitoring and management strategies of ecosystems in biosphere reserves and other conservation areas. By grasping the factors steering vegetation dynamics and habitat alterations, conservation endeavors can be more well-informed and customized to uphold biodiversity and ecosystem resilience amidst ongoing environmental transformations(Fan et al., 2015).



## 7 References

- Aini Rahmi, M. A., Parikesit, P., & Withaningsih, S. (2024). Vegetation change analysis using Normalized Difference Vegetation Index (NDVI) in Sumedang Regency. *E3S Web of Conferences*, 495. <https://doi.org/10.1051/e3sconf/202449502007>
- Al Shawabkeh, R., AlHaddad, M., Al-Fugara, A., Al-Hawwari, L., Al-Hawwari, M. I., Omoush, A., & Arar, M. (2023). Modeling the impact of urban land cover features and changes on the land surface temperature (LST): The case of Jordan. *Ain Shams Engineering Journal*. <https://doi.org/10.1016/j.asej.2023.102359>
- Al-Doski, J., Mansor, S. B., Zulhaidi, H., & Shafri, M. (n.d.). NDVI Differencing and Post-classification to Detect Vegetation Changes in Halabja City, Iraq. In *IOSR Journal of Applied Geology and Geophysics* (Vol. 1, Issue 2). [www.iosrjournals.org](http://www.iosrjournals.org)
- Almalki, R., Khaki, M., Saco, P. M., & Rodriguez, J. F. (2022). Monitoring and Mapping Vegetation Cover Changes in Arid and Semi-Arid Areas Using Remote Sensing Technology: A Review. *Remote Sensing 2022*, Vol. 14, Page 5143, 14(20), 5143. <https://doi.org/10.3390/RS14205143>
- Aslami, F., & Ghorbani, A. (2018). Object-based land-use/land-cover change detection using Landsat imagery: a case study of Ardabil, Namin, and Nir counties in northwest Iran. *Environmental Monitoring and Assessment*, 190(7). <https://doi.org/10.1007/s10661-018-6751-y>
- Badamasi, \*, Yelwa, Abdulrahim, M. A., & Noma. (n.d.). *NDVI threshold classification and change detection of vegetation cover at the Falgore Game Reserve in Kano State, Nigeria*.
- Camps-Valls, G., Campos-Taberner, M., Moreno-Martínez, Á., Walther, S., Duveiller, G., Cescatti, A., Mahecha, M. D., Muñoz-Marí, J., García-Haro, F. J., Guanter, L., Jung, M., Gamon, J. A., Reichstein, M., & Running, S. W. (2021). A unified vegetation index for quantifying the terrestrial biosphere. In *Sci. Adv* (Vol. 7). <https://www.science.org>
- Chen, Y., Wang, K., Lin, Y., Shi, W., Song, Y., & He, X. (2015). Balancing green and grain trade. *Nature Geoscience*, 8(10), 739–741. <https://doi.org/10.1038/NGEO2544>
- Cong, N., Wang, T., Nan, H., Ma, Y., Wang, X., Myneni, R. B., & Piao, S. (2013). Changes in satellite-derived spring vegetation green-up date and its linkage to climate in China from 1982 to 2010: A multimethod analysis. *Global Change Biology*, 19(3), 881–891. <https://doi.org/10.1111/GCB.12077>
- de la Fuente, A., & Williams, S. E. (2022). Climate change threatens the future of rain forest ringtail possums by 2050. *Diversity and Distributions*. <https://doi.org/10.1111/ddi.13652>
- Ding, Y., Xu, J., Wang, X., Peng, X., & Cai, H. (2020). Spatial and temporal effects of drought on Chinese vegetation under different coverage levels. *Science of the Total Environment*, 716. <https://doi.org/10.1016/J.SCITOTENV.2020.137166>

- Dubovyk, O., Landmann, T., Dietz, A., & Menz, G. (2016). Quantifying the impacts of environmental factors on vegetation dynamics over climatic and management gradients of central asia. *Remote Sensing*, 8(7). <https://doi.org/10.3390/RS8070600>
- Ettehadi Osgouei, P., & Kaya, S. (2017). Analysis of land cover/use changes using Landsat 5 TM data and indices. *Environmental Monitoring and Assessment*, 189(4). <https://doi.org/10.1007/s10661-017-5818-5>
- Fan, H., Fu, X., Zhang, Z., & Wu, Q. (2015). Phenology-based vegetation index differencing for mapping of rubber plantations using landsat OLI data. *Remote Sensing*, 7(5), 6041–6058. <https://doi.org/10.3390/rs70506041>
- Filippi, A., Braidot, E., Petrussa, E., Fabro, M., Vuerich, M., & Boscutti, F. (2021). Plant growth shapes the effects of elevation on the content and variability of flavonoids in subalpine bilberry stands. *Plant Biology*, 23(2), 241–249. <https://doi.org/10.1111/plb.13194>
- Fu, B., Wang, Y. K., Xu, P., Yan, K., & Li, M. (2014). Value of ecosystem hydropower service and its impact on the payment for ecosystem services. *Science of the Total Environment*, 472, 338–346. <https://doi.org/10.1016/J.SCITOTENV.2013.11.015>
- Fu, Y. H., Zhou, X., Li, X., Zhang, Y., Geng, X., Hao, F., Zhang, X., Hanninen, H., Guo, Y., & De Boeck, H. J. (2021). Decreasing control of precipitation on grassland spring phenology in temperate China. *Global Ecology and Biogeography*, 30(2), 490–499. <https://doi.org/10.1111/GEB.13234>
- Gado, T. A., El-Hagrasy, R. M., & Rashwan, I. M. H. (2019). Spatial and temporal rainfall changes in Egypt. *Environmental Science and Pollution Research*, 26(27), 28228–28242. <https://doi.org/10.1007/S11356-019-06039-4>
- Han, D., Gao, C., Liu, H., Yu, X., Li, Y., Cong, J., & Wang, G. (2020). Vegetation dynamics and its response to climate change during the past 2000 years along the Amur River Basin, Northeast China. *Ecological Indicators*, 117. <https://doi.org/10.1016/J.ECOLIND.2020.106577>
- Hartoyo, A. P. P., Pamoengkas, P., Mudzaky, R. H., Khairunnisa, S., Ramadhi, A., Munawir, A., Komarudin, K., Hidayati, S., & Sunkar, A. (2022). Estimation of vegetation cover changes using normalized difference vegetation index (NDVI) in Mount Halimun Salak National Park, Indonesia. *IOP Conference Series: Earth and Environmental Science*, 1109(1). <https://doi.org/10.1088/1755-1315/1109/1/012068>
- He, L., Guo, J., Yang, W., Jiang, Q., Chen, L., & Tang, K. (2023). Multifaceted responses of vegetation to average and extreme climate change over global drylands. *Science of the Total Environment*, 858. <https://doi.org/10.1016/J.SCITOTENV.2022.159942>
- Hossain, M. L., & Li, J. (2021). NDVI-based vegetation dynamics and its resistance and resilience to different intensities of climatic events. *Global Ecology and Conservation*, 30. <https://doi.org/10.1016/J.GECCO.2021.E01768>
- Huang, S., Tang, L., Hupy, J. P., Wang, Y., & Shao, G. (2021). A commentary review on the use of normalized difference vegetation index (NDVI) in the era of popular remote

- sensing. In *Journal of Forestry Research* (Vol. 32, Issue 1). Northeast Forestry University. <https://doi.org/10.1007/s11676-020-01155-1>
- Hussain, A., Hussain, I., Ali, S., Ullah, W., Khan, F., Rezaei, A., Ullah, S., Abbas, H., Manzoom, A., Cao, J., & Zhou, J. (2023). Assessment of precipitation extremes and their association with NDVI, monsoon and oceanic indices over Pakistan. *Atmospheric Research*, 292. <https://doi.org/10.1016/j.atmosres.2023.106873>
- Islam, H., Abbasi, H., Karam, A., Chughtai, A. H., & Ahmed Jiskani, M. (2021). Geospatial analysis of wetlands based on land use/land cover dynamics using remote sensing and GIS in Sindh, Pakistan. *Science Progress*, 104(2). <https://doi.org/10.1177/00368504211026143>
- Jiao, K., Gao, J., & Liu, Z. (2021). Precipitation Drives the NDVI Distribution on the Tibetan Plateau While High Warming Rates May Intensify Its Ecological Droughts. *Remote Sensing* 2021, Vol. 13, Page 1305, 13(7), 1305. <https://doi.org/10.3390/RS13071305>
- Johansson, M., Callaghan, T. V., Bosiö, J., Åkerman, H. J., Jackowicz-Korczynski, M., & Christensen, T. R. (2013). Rapid responses of permafrost and vegetation to experimentally increased snow cover in sub-arctic Sweden. *Environmental Research Letters*, 8(3). <https://doi.org/10.1088/1748-9326/8/3/035025>
- Khalile, L., Rhinane, H., Kaoukaya, A., & Lahlaoui, H. (2018). Forest Cover Monitoring and Change Detection in Nfifikh Forest (Morocco). *Journal of Geographic Information System*, 10(02), 219–233. <https://doi.org/10.4236/jgis.2018.102011>
- Knapp, A. K., & Smith, M. D. (2001). Variation among biomes in temporal dynamics of aboveground primary production. *Science*, 291(5503), 481–484. <https://doi.org/10.1126/SCIENCE.291.5503.481>
- Lambin, E. F., Turner, B. L., Geist, H. J., Agbola, S. B., Angelsen, A., Bruce, J. W., Coomes, O. T., Dirzo, R., Unther Fischer, G. U., Folke, C., George, P. S., Homewood, K., Imbernon, J., Leemans, R., Li, X., Moran, E. F., Mortimore, M., Ramakrishnan, P. S., Richards, J. F., ... Xu, J. (2001). The causes of land-use and land-cover change: moving beyond the myths. In *Global Environmental Change* (Vol. 11).
- Li, P., Zhu, D., Wang, Y., & Liu, D. (2020). *Elevation dependence of drought legacy effects on vegetation greenness over the Tibetan Plateau*.
- Li, S., Xu, L., Chen, J., Jiang, Y., Sun, S., Yu, S., Tan, Z., & Li, X. (2023). Monitoring vegetation dynamics (2010–2020) in Shengnongjia Forestry District with cloud-removed MODIS NDVI series by a spatio-temporal reconstruction method. *The Egyptian Journal of Remote Sensing and Space Science*, 26(3), 527–543. <https://doi.org/10.1016/j.ejrs.2023.06.010>
- Li, Y., Li, Z. L., Wu, H., Zhou, C., Liu, X., Leng, P., Yang, P., Wu, W., Tang, R., Shang, G. F., & Ma, L. (2023). Biophysical impacts of earth greening can substantially mitigate regional land surface temperature warming. *Nature Communications*, 14(1). <https://doi.org/10.1038/S41467-023-35799-4>
- Lian, X., Piao, S., Chen, A., Wang, K., Li, X., Buermann, W., Huntingford, C., Peñuelas, J., Xu, H., & Myneni, R. B. (2021). Seasonal biological carryover dominates northern

- vegetation growth. *Nature Communications*, 12(1). <https://doi.org/10.1038/S41467-021-21223-2>
- Liang, S., Zhao, X., Liu, S., Yuan, W., Cheng, X., Xiao, Z., Zhang, X., Liu, Q., Cheng, J., Tang, H., Qu, Y., Bo, Y., Qu, Y., Ren, H., Yu, K., & Townshend, J. (2013). A long-term Global LAnd Surface Satellite (GLASS) data-set for environmental studies. *International Journal of Digital Earth*, 6(SUPPL1), 5–33. <https://doi.org/10.1080/17538947.2013.805262>
- Lin, L. C., Huang, P. H., & Weng, L. J. (2017). Selecting Path Models in SEM: A Comparison of Model Selection Criteria. *Structural Equation Modeling*, 24(6), 855–869. <https://doi.org/10.1080/10705511.2017.1363652>
- Lioubimtseva, E., & Henebry, G. M. (2009). Climate and environmental change in arid Central Asia: Impacts, vulnerability, and adaptations. *Journal of Arid Environments*, 73(11), 963–977. <https://doi.org/10.1016/j.jaridenv.2009.04.022>
- Liu, C., Yan, X., & Jiang, F. (2021). Desert vegetation responses to the temporal distribution patterns of precipitation across the northern Xinjiang, China. *Catena*, 206. <https://doi.org/10.1016/j.catena.2021.105544>
- Liu, Q., Piao, S., Janssens, I. A., Fu, Y., Peng, S., Lian, X., Ciais, P., Myneni, R. B., Peñuelas, J., & Wang, T. (2018). Extension of the growing season increases vegetation exposure to frost. *Nature Communications*, 9(1). <https://doi.org/10.1038/S41467-017-02690-Y>
- Liu, Y., Tian, J., Liu, R., & Ding, L. (2021). Influences of Climate Change and Human Activities on NDVI Changes in China. *Remote Sensing 2021, Vol. 13, Page 4326*, 13(21), 4326. <https://doi.org/10.3390/RS13214326>
- Lu, D., Mausel, P., Brondizio, E., & Moran, E. (2004). Change detection techniques. *International Journal of Remote Sensing*, 25(12), 2365–2401. <https://doi.org/10.1080/0143116031000139863>
- Mao, D., Wang, Z., Luo, L., & Ren, C. (2012). Integrating AVHRR and MODIS data to monitor NDVI changes and their: Relationships with climatic parameters in Northeast China. *International Journal of Applied Earth Observation and Geoinformation*, 18(1), 528–536. <https://doi.org/10.1016/J.JAG.2011.10.007>
- Matsushita, B., Yang, W., Chen, J., Onda, Y., & Qiu, G. (2007). Sensitivity of the Enhanced Vegetation Index (EVI) and Normalized Difference Vegetation Index (NDVI) to Topographic Effects: A Case Study in High-Density Cypress Forest. *Sensors*, 7, 2636–2651. [www.mdpi.org/sensors](http://www.mdpi.org/sensors)
- Menzel, A., Sparks, T. H., Estrella, N., Koch, E., Aaasa, A., Ahas, R., Alm-Kübler, K., Bissolli, P., Braslavská, O., Briede, A., Chmielewski, F. M., Crepinsek, Z., Curnel, Y., Dahl, Å., Defila, C., Donnelly, A., Filella, Y., Jatzak, K., Måge, F., ... Züst, A. (2006). European phenological response to climate change matches the warming pattern. *Global Change Biology*, 12(10), 1969–1976. <https://doi.org/10.1111/J.1365-2486.2006.01193.X>
- Nasiri, V., Deljouei, A., Moradi, F., Sadeghi, S. M. M., & Borz, S. A. (2022). Land Use and Land Cover Mapping Using Sentinel-2, Landsat-8 Satellite Images, and Google Earth

- Engine: A Comparison of Two Composition Methods. *Remote Sensing*, 14(9).  
<https://doi.org/10.3390/rs14091977>
- Nemani, R. R., Running, S. W., Pielke, R. A., & Chase, T. N. (1996). Global vegetation cover changes from coarse resolution satellite data. *Journal of Geophysical Research Atmospheres*, 101(D3), 7157–7162. <https://doi.org/10.1029/95JD02138>
- Novillo, C. J., Arrogante-Funes, P., & Romero-Calcerrada, R. (2019). Recent NDVI trends in mainland Spain: Land-cover and phytoclimatic-type implications. *ISPRS International Journal of Geo-Information*, 8(1). <https://doi.org/10.3390/IJGI8010043>
- Papagiannopoulou, C., Miralles, D. G., Dorigo, W. A., Verhoest, N. E. C., Depoorter, M., & Waegeman, W. (2017). Vegetation anomalies caused by antecedent precipitation in most of the world. *Environmental Research Letters*, 12(7). <https://doi.org/10.1088/1748-9326/aa7145>
- Pei, F., Zhou, Y., & Xia, Y. (2021). Application of normalized difference vegetation index (NDVI) for the detection of extreme precipitation change. *Forests*, 12(5).  
<https://doi.org/10.3390/F12050594>
- Piao, S., Fang, J., Zhou, L., Ciais, P., & Zhu, B. (2006). Variations in satellite-derived phenology in China's temperate vegetation. *Global Change Biology*, 12(4), 672–685.  
<https://doi.org/10.1111/J.1365-2486.2006.01123.X>
- Poyatos, Rafael, L., & Pilar. (n.d.). *Land Use and Land Cover Change After Agricultural Abandonment*. <https://doi.org/10.1659/0276>
- Propastin, P. A., Kappas, M., & Muratova, N. R. (2008). A remote sensing based monitoring system for discrimination between climate and human-induced vegetation change in Central Asia. *Management of Environmental Quality: An International Journal*, 19(5), 579–596. <https://doi.org/10.1108/14777830810894256>
- Rechid, D., Raddatz, T. J., & Jacob, D. (2009). Parameterization of snow-free land surface albedo as a function of vegetation phenology based on MODIS data and applied in climate modelling. *Theoretical and Applied Climatology*, 95(3–4), 245–255.  
<https://doi.org/10.1007/S00704-008-0003-Y>
- Rutkoski, J., Poland, J., Mondal, S., Autrique, E., Pérez, L. G., Crossa, J., Reynolds, M., & Singh, R. (2016). Canopy temperature and vegetation indices from high-throughput phenotyping improve accuracy of pedigree and genomic selection for grain yield in wheat. *G3: Genes, Genomes, Genetics*, 6(9), 2799–2808.  
<https://doi.org/10.1534/g3.116.032888>
- Seo, H., & Kim, Y. (2021). Role of remotely sensed leaf area index assimilation in eco-hydrologic processes in different ecosystems over East Asia with Community Land Model version 4.5 – Biogeochemistry. *Journal of Hydrology*, 594.  
<https://doi.org/10.1016/J.JHYDROL.2021.125957>
- Suepa, T., Qi, J., Lawawirojwong, S., & Messina, J. P. (2016). Understanding spatio-temporal variation of vegetation phenology and rainfall seasonality in the monsoon Southeast Asia. *Environmental Research*, 147, 621–629.  
<https://doi.org/10.1016/J.ENVRES.2016.02.005>

- Tang, X., Li, H., Griffiths, T. J., Xu, X., Ding, Z., & Liu, G. (2015). Tracking ecosystem water use efficiency of cropland by exclusive use of MODIS EVI data. *Remote Sensing*, 7(9), 11016–11035. <https://doi.org/10.3390/RS70911016>
- Tao, J., Xu, T., Dong, J., Yu, X., Jiang, Y., Zhang, Y., Huang, K., Zhu, J., Dong, J., Xu, Y., & Wang, S. (2018). Elevation-dependent effects of climate change on vegetation greenness in the high mountains of southwest China during 1982–2013. *International Journal of Climatology*, 38(4), 2029–2038. <https://doi.org/10.1002/JOC.5314>
- Tu, C., He, T., Lu, X., Luo, Y., & Smith, P. (2018). Extent to which pH and topographic factors control soil organic carbon level in dry farming cropland soils of the mountainous region of Southwest China. *Catena*, 163, 204–209. <https://doi.org/10.1016/J.CATENA.2017.12.028>
- Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A multiscalar drought index sensitive to global warming: The standardized precipitation evapotranspiration index. *Journal of Climate*, 23(7), 1696–1718. <https://doi.org/10.1175/2009JCLI2909.1>
- Yu, L., Xue, Y., & Diallo, I. (2021). Vegetation greening in China and its effect on summer regional climate. *Science Bulletin*, 66(1), 13–17. <https://doi.org/10.1016/J.SCIB.2020.09.003>
- Zhang, P., Cai, Y., Yang, W., Yi, Y., Yang, Z., & Fu, Q. (2020). Contributions of climatic and anthropogenic drivers to vegetation dynamics indicated by NDVI in a large dam-reservoir-river system. *Journal of Cleaner Production*, 256. <https://doi.org/10.1016/J.JCLEPRO.2020.120477>
- Zhang, X., Friedl, M. A., Schaaf, C. B., & Strahler, A. H. (2004). Climate controls on vegetation phenological patterns in northern mid- and high latitudes inferred from MODIS data. *Global Change Biology*, 10(7), 1133–1145. <https://doi.org/10.1111/J.1529-8817.2003.00784.X>
- Zhang, Y., Qi, W., Zhou, C., Ding, M., Liu, L., Gao, J., Bai, W., Wang, Z., & Zheng, D. (2014). Spatial and temporal variability in the net primary production of alpine grassland on the Tibetan Plateau since 1982. *Journal of Geographical Sciences*, 24(2), 269–287. <https://doi.org/10.1007/S11442-014-1087-1>
- Zheng, K., Tan, L., Sun, Y., Wu, Y., Duan, Z., Xu, Y., & Gao, C. (2021). Impacts of climate change and anthropogenic activities on vegetation change: Evidence from typical areas in China. *Ecological Indicators*, 126. <https://doi.org/10.1016/J.ECOLIND.2021.107648>
- Zheng, K., Wei, J. Z., Pei, J. Y., Cheng, H., Zhang, X. L., Huang, F. Q., Li, F. M., & Ye, J. S. (2019). Impacts of climate change and human activities on grassland vegetation variation in the Chinese Loess Plateau. *Science of The Total Environment*, 660, 236–244. <https://doi.org/10.1016/J.SCITOTENV.2019.01.022>
- Zhou, B., Gu, L., Ding, Y., Shao, L., Wu, Z., Yang, X., Li, C., Li, Z., Wang, X., Cao, Y., Zeng, B., Yu, M., Wang, M., Wang, S., Sun, H., Duan, A., An, Y., Wang, X., &

Kong, W. (2011). The Great 2008 Chinese Ice Storm Its socioeconomic-ecological impact and Sustainability Lessons Learned. *Bulletin of the American Meteorological Society*, 92(1),